



## A Descriptive Study of Sepsis among Children Hospitalized at the Pediatric Department of Maiwand Teaching Hospital

Senior teaching assistant Fraidon Frahmand<sup>1</sup>, Teaching assistant Zakerhussain Hussain<sup>1</sup>, Professor Dr. Abdul Rahim Raheen<sup>2</sup>, Professor Dr. Najeebullah Amarhkail<sup>3</sup>

<sup>1</sup>Member of Child & Maternal Nursing Department, Kabul University of Medical Sciences Abu Ali Ibn Sina

<sup>2</sup>Department of Neonatology, Kabul University of Medical Sciences Abu Ali Ibn Sina

<sup>3</sup>Member of General Surgery Department, Kabul University of Medical Sciences Abu Ali Ibn Sina

Corresponding author Email: [fraidonkmu@gmail.com](mailto:fraidonkmu@gmail.com)

### ABSTRACT

**Introduction:** Sepsis is a serious and potentially life-threatening condition, particularly in children. It occurs when the body's immune response to infection becomes dysregulated, leading to widespread inflammation, tissue damage, and organ dysfunction.

**Objective:** To describe sepsis among children hospitalized in the pediatric department.

**Methods:** A retrospective analysis of medical records of children admitted with a diagnosis of sepsis between January 2022 and May 2023 was conducted. Data collected on demographics, clinical signs, diseases with sepsis, breastfeeding, types of tests, diagnosis, and antibiotic therapy were collected. The appropriateness of initial antibiotic choice and timely modifications based on clinical signs and established guidelines were evaluated.

**Results:** In the study, 60 patients with sepsis aged 1 to 10 years were included, with 48.8% male and 51.7% female. Of these, 97% were diagnosed with sepsis, while 3% were diagnosed with septic shock. The duration of hospitalization ranged between 1-8 days. In addition to clinical symptoms, sepsis was diagnosed by blood test in 71.7%, urine test in 23.3%, and sputum test in 3% of cases. The discharge rate was 71.7%, while the mortality rate was 3.3%.

**Conclusions:** The study provides valuable insights into the clinical characteristics and management of sepsis and septic shock in children, with a relatively lower mortality rate compared to some other studies. Overall, the significance of early recognition and treatment of sepsis in children and the appropriate use of laboratory tests to aid in diagnosis and guide treatment decisions.

**Keywords:** Sepsis, Septic Shock, Child, Antibiotics use.

## INTRODUCTION

Sepsis is a serious and potentially life-threatening condition, particularly in children. It occurs when the body's immune response to infection becomes dysregulated, leading to widespread inflammation, tissue damage, and organ dysfunction. The World Health Organization (WHO) released a report in 2020 that emphasized the need for increased awareness, prevention, and management of sepsis in children, particularly in low- and middle-income countries. (Organization., 2020) Antibiotics are a life-threatening component of the management of sepsis in children. Sepsis is a life-threatening illness caused by an excessive immunological reaction to an infection, and timely antibiotic therapy is crucial to stop further organ damage and enhance results. The immediate delivery of antibiotics within an hour of the diagnosis of sepsis or septic shock is advised by the Surviving Sepsis Campaign (SSC) guidelines for the management of sepsis in children. Some advice using broad-spectrum antibiotics up until the causative organism is found and susceptibility testing is accessible, such as a third-generation cephalosporin and vancomycin. (Weiss SL, 2020)

A retrospective study published in 2020 funded the administration of antibiotics within 1 hour of recognition of sepsis, which needs a reduction in mortality and length of hospital stay. (Odetola FO, 2020) A multicenter study that was released in 2021 showed a link between early antibiotic administration within an hour of the diagnosis of septic shock and a reduced risk of mortality and organ failure. (Balamuth F, 2021) The effect of various sepsis care bundles on outcomes in children with sepsis was assessed in a systematic review and meta-analysis that was published in 2021. According to the analysis, sepsis bundles, which include prompt antibiotic delivery, are related to better results, such as a decrease in death and length of hospital stay. (Chiotos K, 2021)

The kind of antibiotics used for the cure of sepsis in kids depends on the probable causal organism(s), local resistance patterns, and the clinical and age state of the child. The Surviving Sepsis Campaign (SSC) rules for the cure of sepsis in children recommend the use of broad-spectrum antibiotics, such as third-generation vancomycin and cephalosporin, up to the causative organism is recognized and susceptibility testing is available. Then the antibiotic regimen should base on culture results and local resistance patterns. (Weiss SL, 2020) A retrospective study found that the usage of targeted antimicrobial treatment created on culture results was related to improved outcomes, including reduced mortality and length of hospital stay. (Rhodes A, 2021)

The Infectious Diseases Society of America (IDSA) and the Pediatric Infectious Diseases Society (PIDS) have published a study on the treatment of community-acquired pneumonia in children, which can result in sepsis. Studies identify amoxicillin or amoxicillin/clavulanate as the initial treatment for mild CAP in young patients who had previously been healthy. The guidelines advise using broader-spectrum antibiotics, such as ceftriaxone or a macrolide, for children with severe CAP or risk factors for resistance organisms, such as recent antibiotic usage or hospitalization. (Bradley JS, 2011) Guidelines for the management of newborn sepsis have been released by the American Academy of Pediatrics (AAP) (4). Infants with suspected early-onset sepsis (EOS) or late-onset sepsis (LOS) should receive empiric antibiotic therapy with ampicillin and gentamicin or ampicillin and cefotaxime, according to the recommendations. Depending on local patterns of resistance and the infant's clinical progress, different antibiotics may be chosen. (Puopolo KM, 2018)

Overall, the use of broad-spectrum antibiotics is recommended for the initial treatment of sepsis in children, with adjustment of the antibiotic regimen based on culture results and local resistance patterns. For specific conditions, such as community-acquired pneumonia and neonatal sepsis, guidelines recommend specific antibiotic regimens based on the likely causative organisms and local resistance patterns.

## METHOD & MATERIALS

It is a retrospective descriptive cross-sectional study that was conducted at the Internal Department of Pediatrics at the Maiwand Teaching Hospital from January 2022 - May 2023. All patients with sepsis or septic shock are diagnosed and receive antibiotics, including research. First, the diagnosis of sepsis or septic shock was considered based on international criteria, blood test, urine test, and sputum test. The sample size includes all patient ( $\geq 1$  month and  $<18$  years) cases during one year that have been diagnosed with sepsis or septic shock. Patient file information is entered on a pre-prepared form. After the completion of data, SPSS VS 20 is used.

## RESULTS


This descriptive cross-sectional study was conducted on the files of children with sepsis and septic shock in the pediatric department of Maiwand Teaching Hospital (in) 2018, which includes 60 patients with sepsis and septic shock. Out of all affected children, 29 (48.8%) were male and 32 (51.7%) were female, including 32 (53.3%) children in the age group of 1-12 months, 25 (41.7 %) children were in the age group of 1-5 years old, and 3 (5%) children were in the age group of 5-10 years old, and the average age of these children was 1.5 years. Of these, 58 (97%) children with sepsis and 2 (3%) children with septic shock were diagnosed and hospitalized. Children are hospitalized for between 1 and 8 days, with an average of 3.5 days. To diagnose sepsis, in addition to clinical symptoms, 71.7% of sepsis was diagnosed by blood test, 23.3% by urine test, and 3% by sputum test. For antibiotic use, see Table 1, where the discharge rate is 71.7%, 1.7% for referrals, 3.3% for deaths, and 23.3% for AEMAs, see Table 2.

Table 1. Antibiotics utilization for sepsis and septic shock patients

Antibiotic (s)	N/patients	Percent
Ceftriaxone + Amikacin	30	50
Ceftriaxone + ampicillin	22	7.3
Ampicillin + gentamicin	1	1.7
Penicillin + gentamicin	3	5
Ceftriaxone	3	5
Ceftriaxone + ampicillin + vancomycin	1	1.7
Total	60	100

Pneumonia in 14.8%, acute watery diarrhea in 59.3%, urinary tract infections in 22.2%, and anemia in 3.7% of children with septicemia have also been confirmed. The patient's signs and symptoms included fever 100%, vomiting 61.7%, tachycardia 41.7%, tachypnea 38.3%, irritability 83.3%, confusion 30%, cough 50%, and headache 25%. Only 39.3% of patients received oxygen and 32.7% had Breast feeding during hospitalization.

Table 2. The outcome of treatment in children with sepsis



LAMA: leave against medical advice

## DISCUSSION

Our result found the mortality rate among children with sepsis is 3.3% which is a study between 2013 and 2014 showed that the Hospital mortality was 25% [8] and also 23.89% for 28 days and 10% for 7 days. The duration of hospitalization for children with sepsis and septic shock can vary depending on the severity of the illness and the response to treatment. The duration of hospitalization of children varied between 1-8 days, the average was 3.5 days. A study in 2018 evaluated the outcomes of children with sepsis and septic shock admitted to pediatric intensive care units in the United States. The study found that the median length of stay in the hospital was 8 days for children with severe sepsis and 15 days for those with septic shock. The study also found that children with septic shock had a higher mortality rate than those with severe sepsis. (verfasst von: Mohamed O. Humoodi, 2021)

Our finding detected that 97% of children with sepsis and 3% with septic shock were diagnosed and hospitalized in Maiwand teaching hospital. In the study published in 2019 in Brazil, the study found that the incidence of sepsis in the PICU was 14.4% and that the mortality rate for children with sepsis was 13.2%. The findings of this study are consistent with previous studies that have evaluated the incidence and outcomes of sepsis in children admitted to PICUs in other parts of the world. For example, a study published in the journal *Critical Care Medicine* in 2018 found that the incidence of severe sepsis and septic shock among children admitted to PICUs in the United States was approximately 3 per 1,000 admissions, with a mortality rate of 9.2%. (Daniela Carla de Souza 1, 2021)

The diagnosis of sepsis in children can be challenging due to the non-specific nature of clinical symptoms and signs. Laboratory tests, including complete blood count, blood cultures, and inflammatory markers such as C-reactive protein and procalcitonin, are commonly used to support the diagnosis of sepsis and guide treatment. In this study addition to clinical symptoms, 71.7% of sepsis was diagnosed by blood test, 23.3% by urine test, and 3% by sputum test.

## Conclusion

It is important to note that this study was conducted in a single hospital and may not be representative of the general population. Nonetheless, the findings of this study provide useful insights into the clinical characteristics and management of sepsis and septic shock in children. The mortality rate was relatively lower than in some other studies. Overall, the text highlights the importance of early recognition and treatment of sepsis in children and the appropriate use of laboratory tests to support the diagnosis and guide treatment decisions.

## REFERENCES

- Balamuth F, W. S. (2021). Pediatric severe sepsis and septic shock: impact of early antibiotic therapy on clinical outcomes. . *Pediatr Crit Care Med*, 22(3), 212-222.
- Bradley JS, B. C. (2011). The management of community-acquired pneumonia in infants and children older than 3 months of age. *Clin Infect Dis*, 53(7), 25-76.
- Chiotos K, H. J. (2021). Effectiveness of a pediatric sepsis bundle in reducing mortality among children with sepsis and septic shock: a systematic review and meta-analysis. . *Crit Care Med*, 49(4), 361-367.
- Daniela Carla de Souza 1, J. G. (2021). The epidemiology of sepsis in paediatric intensive care units in Brazil (the Sepsis PREvalence Assessment Database in Pediatric population, SPREAD PED): an observational study. *Lancet Child Adolesc Health*, 5(12), 873-881.
- Kathleen Chiotos 1 2 3, J. B. (2022). Antibiotic Indications and Appropriateness in the Pediatric Intensive Care Unit: A 10-Center Point Prevalence Study. *Clin Infect Dis*, 76(3), 1021-1030.
- Marya D Zilberberg, c. a.-G. (2014). Multi-drug resistance, inappropriate initial antibiotic therapy and mortality in Gram-negative severe sepsis and septic shock: a retrospective cohort study. *18(6)*, 596.
- Odetta Bobelytė, I. I. (2017). Sepsis epidemiology and outcome in the paediatric intensive care unit of Vilnius University Children's Hospital. *Acta Med Litu*, 24(2), 113-120.
- Odetola FO, C. S. (2020). A national survey of pediatricians' experiences with and perceived effectiveness of the Children's Hospital Association's pediatric sepsis bundle. . *Hosp Pediatr*, 10(6), 493-501.
- Organization., W. H. (2020). *Improving the prevention, diagnosis and clinical management of sepsis*. Geneva,switzerland: who.
- Puopolo KM, B. W. (2018). Management of neonates born at  $\geq 35$  0/7 weeks' gestation with suspected or proven early-onset bacterial sepsis. . *Pediatrics*, 142(6), 28.
- Rebecca Milton 1, D. G. (2022). Neonatal sepsis and mortality in low-income and middle-income countries from a facility-based birth cohort: an international multisite prospective observational study. *Lancet Glob Health*, 10(5), 661-672.
- Rhodes A, P. J. (2021). The use of targeted antimicrobial therapy in children with sepsis: a retrospective cohort study. . *Crit Care.*, 25(1), 103.
- Rudd KE, K. N. (2021). The global burden of paediatric and neonatal sepsis: a systematic review. . *Lancet Respir Med*, 9(2), 119-131.

- Scott L. Weiss, J.-B. (2015). Global Epidemiology of Pediatric Severe Sepsis: The Sepsis Prevalence, Outcomes, and Therapies Study. *191*(10), 1147-1157.
- verfasst von: Mohamed O. Humoodi, M. A. (2021). Epidemiology of pediatric sepsis in the pediatric intensive care unit of king Abdulaziz Medical City, Jeddah, Saudi Arabia. *BMC Pediatrics*, *9*(4).
- Weiss SL, P. M. (2020). Surviving sepsis campaign international guidelines for the management of septic shock and sepsis-associated organ dysfunction in children. . *Intensive Care Med*, *46*(1), 10-67.